


Theme 3: Fodder management and conservation

GUIDELINES TO MAIZE SILAGE MAKING



A publication sponsored by the ICSTIAPL project





You will learn about (learning objectives):

- The rights steps for making maize silage
- Identifying the right stage of harvesting, using the right machinery for best results and ensuring proper management of pit/clamp before and after feed out.

Making Maize silage

Maize silage is a way of preserving maize fodder for future use

The silage provides low-cost metabolizable energy (ME) for dairy cows and can be a key ingredient in a cow's daily feed ration/diet

Maize silage is low in Crude protein (CP), Calcium and Vitamin D

If prepared in enough quantity, it helps a farmer to counter seasonality in milk production



Stage of harvesting maize fodder for silage

The ideal stage of harvesting is when the kernel is dough-ripe / ripe

That is, when twisting the stem, only few drops of moisture can still be seen coming out – see next slide

The leaves below the cob are dry

The milk line is at half to three quarters of the kernel

Aim at a DM level of 30-35% for the whole crop



Estimating the dry matter on the stem

Take the maize stem, twist it on your knee as shown alongside

- 4 drops \leq 18 % DM 100 % green
- 3 drops \approx 21 % DM 75 % green
- 2 drops \approx 24 % DM 50 % green
- 1 drop \approx 27 % DM 25 % green
- 0 drops \geq 30 % DM 0 % green



Chopping

Chop length should be between 8 – 12mm

Sharpen the knives at least every day

When harvesting is done at dough-ripe stage the chopper needs a kernel crusher

Preventive maintenance and servicing of the machines during harvesting should be done daily



Location of the pit

The location of the silo (silage pit) should;

- be dry with no stagnant water
- be protected from animals and wildlife
- have a high elevation or on a slope
- be as close as possible to the cow or cow barn.



Transportation

The choice of means of transport is determined by the amount of harvested maize, distance and road network

The distance between the field and pit/clump must not be more than 3km, if motorized transport is available

Plan in such a way that silage pit can be covered in 12 hours



Filling of the Silo: Pit/clamp / bag

Fill the pit layer by layer. Make layers of 10 cm silage every time

Additives such as molasses can be added at this stage

Compact after every layer

Prevent contamination with soil, dirt and manure.



Silage pit dimensions and location

The silage pit should be narrow and long

Determine how far the clamp/pit is from the barn, how well-drained the location is, how safe from any other traffic and from birds, rodents and wild animals.



Compacting

Compaction is necessary to remove air (oxygen) from the chopped maize

Tools used for compaction need to have enough weight for effective compaction

Ensure equal compaction of the silage surface.



Sealing

Ensure that the polythene sheet is not punctured and has a good overlap to cover the silage pit / clamp completely

Ensure that the thickness of polythene is >500 gauge

Seal silage pits or clamps within 24 hours after starting to make the pit

Care needs to be taken that the polythene is not damaged.



Covering with soil

Soil cover of at least 20-30cm is important for four reasons:

1. Continuously compacts the silage
2. Protects the polythene sheet against direct sunlight
3. Protect the polythene sheet from being damaged by birds and small wildlife
4. Protect the silage underneath the polythene sheet from heating up.

Soil or sandbags keep the polythene sheet tighter and continue exerting pressure on the silage pit

Tires are not preferred; they do not keep out the heat. Tires can also puncture polythene.



Silage making in transportable bales

Baling eases transportation of the ready-made silage from one location to another

This can be useful for farmers with dairy cows and not enough land to plant their own fodder crops

Also, farmers in dry or very dry areas where rainfall pattern does not support the growth of forages can benefit from this technology.



Fermentation

Sometimes circumstances for ensiling may not be very ideal, for example; the maize is too wet or too dry. To enhance fermentation, additives may be added during filling of the pit

Ensure regular and equal application of the additive over the silage surface

After closing the pit / clamp, it should stay closed for 6 weeks before opening



Making a narrow pit / clamp will ensure feeding speed on feed out.

Feed Out

The feeding speed should be 1.5-2meters per week(20-30cm per day)

Slow feeding speed will increase the risk of secondary fermentation.

Make sure that all the silage is removed from the face of the pit to avoid heating up.

Remove all loose materials from the open side (face) of the pit and feed immediately. Loose materials in front or on the bottom of the silage pit or clamp cause moulds and rotting bacteria to grow. From here moulding and rotting will spread to the good parts of the clamp

Clean the open silage face from all rotten and loose materials daily. This removes moulding and rotting materials therefore preventing contamination of good feed.



Feeding

A mature cow can consume between 15-35 kg of maize silage per day

Feed more maize silage in the early lactation period when the cows milk production is highest

Maize silage contains very little protein, supplement milking cows with high protein feeds.







Ministry of Foreign Affairs of the
Netherlands

About the ICSIAPL Project

The Integrated & Climate Smart Innovations for Agro-Pastoralist Economies and Landscapes in Kenya's Arid and Semi-arid Lands (ICSIAPL) is a three-year (2021 -2023) project funded by the European Union (EU) and the Ministry of Foreign Affairs of the Kingdom of the Netherlands (DGIS). It is managed through a delegated cooperation with the Embassy of the Kingdom of the Netherlands (EKN) in Nairobi.

The EU fund is attached to the DeSIRA Initiative (Development Smart Innovation through Research in Agriculture).

The project is implemented by SNV Netherlands Development Organisation, SNV, (lead partner) and the Kenya Agriculture Livestock Research Organization, KALRO (research partner).

